

Identical mutation in a novel retinal gene causes progressive rod–cone degeneration in dogs and retinitis pigmentosa in humans

Barbara Zangerl^a, Orly Goldstein^b, Alisdair R. Philp^c, Sarah J.P. Lindauer^a,
Susan E. Pearce-Kelling^b, Robert F. Mullins^c, Alexander S. Graphodatsky^d, Daniel Ripoll^e,
Jeanette S. Felix^f, Edwin M. Stone^{c,g}, Gregory M. Acland^b, Gustavo D. Aguirre^{a,*}

^a *Clinical Studies–Philadelphia, School of Veterinary Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA*

^b *James A. Baker Institute, College of Veterinary Medicine, Cornell University, Ithaca, NY 14853, USA*

^c *Ophthalmology and Visual Sciences, Carver College of Medicine at The University of Iowa, Iowa City, IA 52242, USA*

^d *Institute of Cytology and Genetics, Russian Academy of Sciences, Novosibirsk, Russian Federation*

^e *Computational Biology Service Unit, Cornell Theory Center, Cornell University, Ithaca, NY 14853, USA*

^f *OptiGen, LLC, Cornell Business & Technology Park, Ithaca, NY 14850, USA*

^g *Howard Hughes Medical Institute, Iowa City, IA 52242, USA*

Received 30 May 2006; accepted 13 July 2006

Available online 30 August 2006

Abstract

Progressive rod–cone degeneration (*prcd*) is a late-onset, autosomal recessive photoreceptor degeneration of dogs and a homolog for some forms of human retinitis pigmentosa (RP). Previously, the disease-relevant interval was reduced to a 106-kb region on CFA9, and a common phenotype-specific haplotype was identified in all affected dogs from several different breeds and breed varieties. Screening of a canine retinal EST library identified partial cDNAs for novel candidate genes in the disease-relevant interval. The complete cDNA of one of these, *PRCD*, was cloned in dog, human, and mouse. The gene codes for a 54-amino-acid (aa) protein in dog and human and a 53-aa protein in the mouse; the first 24 aa, coded for by exon 1, are highly conserved in 14 vertebrate species. A homozygous mutation (TGC → TAC) in the second codon shows complete concordance with the disorder in 18 different dog breeds/breed varieties tested. The same homozygous mutation was identified in a human patient from Bangladesh with autosomal recessive RP. Expression studies support the predominant expression of this gene in the retina, with equal expression in the retinal pigment epithelium, photoreceptor, and ganglion cell layers. This study provides strong evidence that a mutation in the novel gene *PRCD* is the cause of autosomal recessive retinal degeneration in both dogs and humans.

© 2006 Elsevier Inc. All rights reserved.

Keywords: Dogs; Disease models, animal; Genetic diversity; Genetic linkage; Genetic markers; Genetic predisposition to disease; Genetic variation; Mutation; Retinal degeneration; Retinitis pigmentosa

Progressive rod–cone degeneration (*prcd*) is a canine, late-onset, autosomal recessive photoreceptor degeneration of unknown molecular etiology [1]. This disorder is one of several inherited diseases grouped under the progressive retinal atrophy (PRA) rubric. These serve as animal models for human retinitis pigmentosa (RP) due to shared molecular defects and/or phenotypic similarities (see [2] for review). Together with the *rd* mouse and RCS rat, *prcd* represents one

of the best characterized models of inherited retinal degeneration (see [3–12]). Previously, the disease was mapped to the centromeric end of canine chromosome 9 (CFA9), a region orthologous to the telomeric end of human chromosome 17 (HSA17q22; [13,14]). A physical and radiation hybrid map of the region confirmed the syntenic organization of the homologous chromosomal segments, but demonstrated also that the order of gene clusters, and genes within clusters, differed between human and dog [15].

In a companion paper, we reported the use of linkage disequilibrium (LD) analysis across several dog breeds

* Corresponding author. Fax: +1 215 573 2162.

E-mail address: gda@vet.upenn.edu (G.D. Aguirre).

affected with *prcd* [2,9] to narrow the candidate region and direct the positional cloning of *prcd* [16]. As individual dog breeds represent genetic isolates, this powerful technique reduced the critical interval from ~6.4 Mb to 106 kb and identified a chromosomal haplotype shared by 14 different dog breeds or breed varieties [16]. This suggested that a common ancestral chromosome containing the mutant gene was widely distributed as modern dog breeds developed. The LD region, flanked by *AANAT* and *ST6GalNac1*, is completely contained in a single RPC181 BAC clone (10-M13; [17]; <http://bacpac.chori.org>) and includes three known genes: *CYGB*, *RHBDL*, and *STHM* (*ST6GalNac2*). All the genes in the interval were evaluated and excluded from causative association with disease [16].

In a complementary effort we created a normalized canine retinal EST library that was catalogued by suggested gene homology and chromosomal location [18]. Intended as a means to identify and evaluate positional candidate genes within the *prcd* interval, the combination of sequence and expression data yielded several potential candidate genes in the original disease-relevant interval. The application of this approach on the final shared haplotype identified a mutation in a novel gene of unknown function, referred to as progressive rod–cone degeneration (*PRCD*). In this study, we report the preliminary characterization of *PRCD*. Furthermore, we analyzed samples from 1863 human patients with RP and other inherited retinal disorders, whose diseases had been previously excluded from causal association with genes known to cause retinal degeneration (RetNet—Retinal Information System; <http://www.sph.uth.tmc.edu/RetNet/>), and demonstrate that the identical mutation that causes *prcd* in dogs is homozygously present in a human RP patient.

Results

Identification and characterization of *PRCD*

Identification of new candidate genes from a retinal EST library

Although linkage analysis of *prcd*-informative pedigrees mapped the disease locus to CFA9 [14], the original canine genome sequence draft (<http://www.genome.gov/11007358>; July 2004) suggested a translocation of the centromeric CFA9 region to CFA18. To establish the correct chromosomal position of the disease interval, FISH hybridization was performed with clones isolated from an 8× redundant dog genomic RPC181 BAC library ([17]; <http://bacpac.chori.org>) that are specific for CFA9 (189-B1) or are included in the physical map of the *prcd* interval (429-O8; [15]). Both probes colocalized on the same canine chromosome that was identified as CFA9 by the chromosome-specific probe and G-banding pattern (data not shown). The assignment of the *prcd*-critical interval to CFA9 was confirmed in the new canine genome draft assembly ([19], May 2005 v2.0).

As presented in a companion paper [16], LD mapping identified a common 106-kb candidate region on CFA9 flanked by *AANAT* and *ST6GalNac1*. This interval, and flanking genes, is contained in RPC181 BAC 10-M13 and includes RefSeq genes *AANAT*, *RHBDL6*, *CYGB*, *ST6GalNac1*, and *ST6GalNac2*, all of which were previously excluded from causative association with disease. To complement the positional cloning of *prcd* and begin characterization of the canine retinome, we developed a normalized canine retinal cDNA library [18] containing a total of 6316 retinally expressed clones that were assembled into 3980 unique sequences, termed “contig,” with unique identifier numbers (<http://www.bioinformatics.upenn>.




Fig. 1. Display of the *prcd* candidate region. (A) LD was used to reduce the candidate region to 106 kb [16]. This interval harbors three canine retinal ESTs (contigs 1131, 1365, and 3041). *PRCD* was cloned within orthologous regions in the (B) dog, (C) human, and (D) mouse and showed a high degree of conservation in gene structure.




Fig. 3. (A) Example of a *prcd*-informative ECS pedigree showing cosegregation of haplotypes with the disease. *GRB2*, *AANAT*, and *SEC14L* are outside the candidate region, but a benign *AANAT* SNP is in complete linkage disequilibrium with the disease in this breed [16]. (B) Restriction enzyme test shows complete digestion of the *PRCD* A allele with *RsaI* and of the G allele with *ApaLI*. (C) Assessment of the *PRCD* genotype by pyrosequencing.

the C2Y mutation in the human and canine sequences (see below) does not appear to change the predicted α -helical conformation of the initial 15-aa fragment. *PRCD* could not be identified in nonvertebrate species (e.g., *Drosophila melanogaster*, *Arabidopsis thaliana*, or any published insect genome).

We screened for potential transcription binding sites 1000 bp upstream of the ATG using the consite algorithm [score threshold of 80% (<http://mordor.cgb.ki.se/cgi-bin/CONSITE/consite>)] or manually (CRX, NRL). Overall, 86, 74, and 74 motifs are located on the positive strand only in the dog, human, and mouse sequence and represent binding sites for 32, 36, and 23 different factors, respectively. Thirteen transcription factors are common to all three species (AML-1, ARNT, bZIP910, E74A, Hen-1, Max, NF-Y, n-MYC, Snail, Sox5, Sox17, Thing1-E47, USF). Pair-wise analysis of alignments, however, identified only one Sox17 site, located approximately 50 bp in front of the ATG, and CRX binding sites that were spatially conserved (Supplementary Fig. 1, dark and light gray boxes, respectively). The same comparative analysis suggests potential involvement of CFI-USP and Snail as regulators through binding sites in intron 1 of *PRCD* (data not shown). CRX binding sites were also recognized in the potential 5'UTR of the additional 11 species for which exon 1 sequence is available, strongly suggesting the importance of this regulatory site. A potential binding motif for another retinal regulator, NRL, was found in the putative promoter region of the chicken, but could not be identified in the other species.

Putative TATA boxes are located at position -565 in dog and -593 in human (AATATT; Supplementary Fig. 1); a second box could be present at -723 and -759 in the dog and human, respectively, but is less conserved between species (common motif ATATA), and the human might have a third box at -803 (not shown in Supplementary Fig. 1 alignment). Putative CAAT motifs are observed at -408 in the dog and at -242 and -595 in human. However, none of these transcription initiation sites could be identified in the mouse sequence.

Mutation in PRCD causes retinal degeneration in dog and human

Identification and segregation of the canine PRCD mutation

Retinal cDNA sequence comparison of the canine *PRCD* ORF (Supplementary Table 1, primers 3 and 4) from 8.6-week *prcd*-affected and 10.4-week normal dogs identified a single G to A transition at nucleotide 5 of the coding sequence. This mutation causes a cysteine to tyrosine change (C2Y) at the second amino acid of the protein in the affected animal. The causal association between this mutation and *prcd* was established by: (a) consistent and stable segregation of the mutation in the *prcd* reference colony; (b) segregation of the mutation in well-maintained pedigrees of breeds included in previous *prcd* breeding studies; (c) segregation of the mutation in dog breeds for which the diagnosis of *prcd* was obtained by the investigators (G.D.A., G.M.A.) and pedigrees were followed over several generations; (d) observation of the *prcd* mutation in breeds with a clinical diagnosis of retinal degeneration

compatible with *prcd*; (e) absence of the mutated allele from breeds not segregating the *prcd* phenotype. Details of these studies are presented below:

- (a) We have maintained a reference colony segregating *prcd* for more than 10 generations. The founder chromosomes were from affected toy and miniature poodles (TP, MP [9]). Affected chromosomes originating from Labrador retriever (LR), Portuguese water dog (PWD), Nova Scotia duck tolling retriever (NSDTR), Australian cattle dog (ACD), and English cocker spaniels (ECS) were introduced into the colony subsequently and segregate the same phenotype. The affected haplotype present in this colony has been identified [16] and shows complete concordance with the phenotype. The *prcd* mutation is included in this haplotype described by 98 polymorphisms in Supplementary Table 1 of the companion paper [16] and does not show any discordance from inheritance within the reference colony.
- (b) We also have access to outside pedigrees from these breeds in which *prcd* segregates. These animals were screened for the disease-specific haplotype and canine *PRCD* mutation. Complete concordance between affected phenotype and genotype was confirmed for affected and obligate carriers in ACD, ECS, LR, NSDTR, MP, PWD, and TP using a restriction enzyme analysis (see Figs. 3A and 3B for illustration). In addition, we have selected a small subset of dogs from some breeds (ACD, LR, MP, TP) that had a clinical retinal disease *atypical* for the breed-specific form of retinal degeneration. For example, these included dogs showing clinical signs much earlier or later than the characteristic age of onset of *prcd* in the breed, inflammatory retinopathy, or other clinical disease features not considered characteristic of *prcd*. In all cases, these dogs were found to be heterozygous or homozygous normal at the disease locus. Sequence analysis using a subset of markers from the *prcd* LD region confirmed exclusion of *prcd* and suggested that the clinical disease (s) present in these dogs were geno- or phenocopies.
- (c) Pedigrees were available for four breeds of which three were not included in the breeding studies [9]: American cocker spaniel, American Eskimo (AE [21]), Chesapeake Bay retriever, and Entlebucher mountain dog (EMD [22]). With the exception of one dog, there was complete concordance between the canine *PRCD* mutation and affected and obligate carrier status. The exception was an AE that was homozygous affected (A/A), but clinically normal at 13 years. As the disease in this and some other breeds, e.g., ECS, can occur at a much later age [2,9], definite phenotypic diagnosis will require histologic assessment after death of the animal.
- (d) Selected samples of representative dog breeds that showed a retinal disease that appeared to be clinically compatible with *prcd* were collected by the investigators or provided by colleagues. These included the following breeds: Chinese crested (CC), silky terrier (ST), Finnish lapphund (FL), Kuvasz, Laponian herder, and Swedish

lapphund. The homozygous G → A mutation in canine *PRCD* was found in all five breeds. Further analysis of these breeds is currently in progress.

- (e) To establish that the A allele is not just a rare SNP in the population that has been concentrated in a few selected breeds, high-throughput pyrosequencing analysis (Fig. 3C) was used to screen 1120 samples from unrelated dogs from two groups: (i) 1007 samples from 37 breeds with a non-*prcd* form of retinal degeneration for which dogs of known phenotype were tested; in several of these breeds, the molecular basis of the retinal disease was known and the genotype at the disease locus established (Supplementary Table 2); (ii) 113 samples of unrelated dogs from 54 breeds without known retinal disease (Supplementary Table 3). All 1120 dogs tested from 91 different breeds were homozygous normal (G/G) at nucleotide 5 of canine *PRCD*. Furthermore, 24 red and 6 gray wolves also were tested, and all were homozygous normal (G/G) at the mutation locus.

In summary, the canine *PRCD* mutation has been observed in 18 individual dog breeds or breed varieties (Table 1) to date and was confirmed to cause disease in the homozygous state. The disease-causing allele was never observed in dogs or breeds confirmed not to segregate *prcd*.

Mutation identification and screening in human populations

We have found seven different human *PRCD* sequence variations in a total of 1863 patients, most of whom were clinically diagnosed with RP or allied retinal degenerations (Table 2). Of these, 4 were found in the heterozygous state in the noncoding regions of the gene and appeared unlikely to represent disease-causing variations. One patient with autosomal recessive RP (arRP) showed a G → A transition in the heterozygous state at 88 bp of the coding sequence. This variation would be expected to cause a conservative substitution of methionine for valine at aa 30 of the *PRCD* protein (V30M).

Table 1
Breeds affected by *prcd*

Abbreviation	Breed
ACD	Australian cattle dog
ACS	American cocker spaniel
AE	American Eskimo
CBR	Chesapeake Bay retriever
CC	Chinese crested
ECS	English cocker spaniels
EMD	Entlebucher mountain dog
FL	Finnish lapphund
KU	Kuvasz
LH	Lapponian herder
LR	Labrador retriever
MP	Miniature poodle
NSDTR	Nova Scotia duck tolling retriever
PWD	Portuguese water dog
SL	Swedish lapphund
ST	Silky terrier
STCD	Australian stumpy tail cattle dog
TP	Toy poodle

DNA sequencing of the entire human *PRCD* coding sequence in this patient failed to detect a second sequence variation. The heterozygosity of the variation in this patient, coupled with the conservative nature of the predicted change in the protein, makes it difficult to assert that this variation causes human retinal disease even though the change was not observed in the control population (215 individuals; Table 2).

Another transition (C → T in the 49th nucleotide of the coding sequence, predicting the substitution of cysteine for arginine in codon 17, R17C) was observed in the heterozygous state in one RP patient of Indian origin, but was also observed in the heterozygous state in five members of the Indian/Pakistani control group. Bidirectional DNA sequencing was used to examine the entirety of the human *PRCD* coding sequence in the RP patient with the R17C change, and no other variations were detected. This result, combined with the high frequency of occurrence (5% of all alleles) of the variation in the ethnic control samples, makes it likely that it represents a non-disease-causing ethnic-specific polymorphism.

In contrast, a woman from Bangladesh with arRP was found to harbor a G → A transition in the fifth nucleotide of the human *PRCD* coding sequence, which would be predicted to result in a substitution of tyrosine for cysteine (C2Y) in the second amino acid of the protein. This is the identical mutation at both the DNA and the protein levels that was observed in *prcd*-affected dogs. The patient was 32 years of age when examined at the University of Iowa. She was born in Bangladesh and first noticed difficulty seeing at night as a small child. Her family history was significant for four unaffected siblings, one brother who became blind at a young age, and two other siblings still living in Bangladesh with an uncharacterized vision problem. Her parents were first cousins and had normal vision. The patient's best corrected visual acuity was 20/70 in the right eye and hand motions in the left eye with a manifest refraction of $-3.75 + 1.00 \times 180$ in the right eye and -2.00 sphere in the left eye. Slit-lamp examination revealed clear corneas and anterior chambers that were deep, clear, and quiet. The lenses were clear, and there were no vitreous cells. Analysis of the papillary response revealed a 0.9 log unit relative afferent papillary defect. Goldmann perimetry showed a greater loss of sensitivity in the left eye than in the right (Fig. 4A). Fundus examination revealed optic discs with fairly normal color, but markedly attenuated arterioles (Fig. 4B). There was extensive bone-spicule-like pigmentation in all four quadrants of both eyes, densest at the equator, and this intraretinal pigment was admixed with lighter colored deposits at the level of the retinal pigment epithelium. There were also small patches of geographic atrophy near fixation in both eyes. Ganzfeld electroretinography was performed, but no responses could be recorded from either eye under any stimulus condition.

Expression and function of the *PRCD* gene

Expression profile of canine *PRCD* in normal and mutant animals

Transcription of canine *PRCD* was examined by northern analysis using probes specific for either exon 1 or exons 2–4

Table 2

Human *PRCD* sequence changes observed in the heterozygous (het) or homozygous (hom) state in patients with RP, Leber congenital amaurosis (LCA), cone-rod dystrophy (CRD), or Usher syndrome (USH) and in normal Caucasian (NML) and Indian/Pakistani (I/P NML) control populations

Disorder			RP	LCA	CRD	USH	NML	I/P NML
Number of patients/controls			1241	283	224	115	166	49
Position	Sequence change	aa change						
-106 het	C → T		1	0	0	0	0	0
-40 U het	T → C		1	0	0	0	0	0
5 hom	TGC → TAC	Cys2Tyr	1	0	0	0	0	0
49 het	CGC → TGC	Arg17Cys	1	0	0	0	0	5
88 het	GTG → ATG	Val30Met	1	0	0	0	0	0
*57 U het	C → G		1	0	0	0	0	0
*80 het	G → A		1	0	0	0	0	0

* indicates the nucleotide 3' of the termination codon; U indicates that the nucleotide is within the UTR.

(Fig. 5A). The same level of expression was observed in retinas from normal and *prcd*-affected dogs and showed two distinct retinal transcripts of 2 and 6.5 kb. Experiments with a probe specific for the 3'UTR located in the contig 1131 sequence yielded the same results (data not shown). The same transcripts were observed in RPE/choroid, but the level of expression in other tissues was too low for detection. RT-PCR amplification using primers located in exons 1 and 4 (Supplementary Table 1, primers 1 and 31) showed low levels of canine *PRCD* expression in a variety of tissues (data not shown).

To quantify the expression profile of canine *PRCD*, we performed quantitative real-time PCR (qRT-PCR) on different tissues of a 16-week-old normal dog for two probes: one specific for exon 1 and the other spanning parts of exons 2 and 3. Results are shown for the exon 1 probe only, as no significant differences between the two probe sets were found. Relative expression was calculated against an 18S probe as the difference of cycles leading to amplification. In addition, retinal samples from *prcd*-affected dogs of various ages (10, 16, and 23 weeks; 2 and 3 years) also were analyzed by qRT-PCR. Comparison of canine *PRCD* expression in the normal vs. *prcd*-affected retina showed

no significant changes in expression with age or disease stage (Fig. 5B). Expression in spleen, liver, and pancreas was insufficient for quantification, and no expression was observed in the lung (data not shown). In comparison to normal retina, there was significantly lower expression in all other tissues tested (Fig. 5B); there was ~6.5-fold lower expression in RPE/choroid, 16-fold decrease in kidney, and 100- to 1000-fold decreases in other tissues. Even though expression levels for cerebrum represented a pooled sample of frontal, occipital, and temporal lobes, analysis of individual samples showed comparable low levels of expression (data not shown).

Laser capture microscopy was used in the normal retina to compare canine *PRCD* expression specifically in the RPE, photoreceptor, and ganglion cell layers. Comparable expression in all three layers suggests that this gene has uniform expression in many retinal cell types and is not photoreceptor specific or enriched (Fig. 5C).

Expression of wild-type and mutant human *PRCD* in COS cells

His-tag localization was performed on COS cells transfected with either wild-type or mutant (C2Y) human *PRCD*. Whereas




Fig. 4. Clinical assessment of an arRP patient of Bangladesh birth at age 32. (A) The Goldmann visual field of the right eye shows complete loss of the I2e isopter. The I4e isopter (shown in blue) is limited to less than 5°. There is persistence of a temporal island that is sensitive only to the V4e target (shown in purple). (B) Fundus photograph of the right eye of the patient centered on the inferotemporal vascular arcade. The optic disk is of fairly normal color, but the retinal arterioles are extensively attenuated. Typical bone-spicule-like pigmentation can be seen admixed with pale deposits at the level of retinal pigment epithelium. Small patches of atrophy of the retinal pigment epithelium can be seen adjacent to the fovea (the fovea can be recognized as the region of darker pigmentation to the left of the optic nerve head).




Fig. 5. Expression profile of canine *PRCD*. (A) Northern analysis with probes specific for exon 1 and exons 2–4 of canine *PRCD* identifying two distinct transcripts (2 and 6.5 kb) expressed in the normal and *prcd*-affected retinas (R, 10.4 weeks; R *prcd*, 8.6 weeks). No signal is present in normal or affected brain (B, 15.7 weeks; B *prcd*, 7.7 weeks) or normal spleen (S, 22.1 weeks). The same transcripts are expressed at lower levels in the RPE/choroid (RPE) compared to retina. β -Actin was used as loading control. (B) qRT-PCR results show high expression in the retina compared to other tissues. An expression profile (shown as relative expression against 18S) was established for a normal dog at 16 weeks of age (center) and is similar to the expression levels found in affected retinas of different ages (left). Relative expression normalized against retina provides fold reduction in the expression between tissues (right). (C) Individual retinal layers (RPE, photoreceptor, and ganglion cell layers) from a normal eye were isolated with a laser capture microscope (respective regions are illustrated by octagons below black bars) and analyzed by qRT-PCR. No significant differences in expression were observed between the layers.

the wild-type fusion protein was distributed diffusely throughout the cytoplasm (Figs. 6A and D), in the majority of transfected cells the mutant protein was present in spherical aggregates within the cytoplasm (Figs. 6E and 6H). The distribution of the wild-type protein did not substantially overlap with that of actin filaments in the cytoplasm, but did extend into cell processes and showed greatest labeling around the nucleus.

Discussion

A combination of LD mapping [16] and the characterization the canine retinome [18] identified the 3' end of a novel retinal

gene. This gene was subsequently cloned in dog, human, and mouse, and a mutation that causes canine and human retinal degeneration was found. The gene has been given the preliminary name of “progressive rod–cone degeneration” with the gene symbol “*PRCD*” (approved by the HUGO Gene Nomenclature Committee).

PRCD has a predominant transcript of 671 bp in the dog retina and encodes a conserved ORF (78–81% conservation). GenBank entries of human and murine transcripts, as well as potential alternative splicing identified in the dog, suggest alternative splice variants for this gene, and some of these have been characterized in dogs (data not shown). However, we could not identify usage of additional 5' exons in any of the




Fig. 6. Immunocytochemical localization of (A) wild-type and (E) mutant *PRCD* in COS-7 cells. No colocalization was observed with phalloidin-labeled actin (B, F) or nuclei stained with DAPI (C, G) in the merged images (D, H). Localization of the His tag was present in a diffuse weblike distribution in most cells transfected with the wild-type construct (A, D), whereas labeling was generally concentrated in vesicular profiles in the cytoplasm in cells transfected with the mutant construct (E, H). Magnification: 460 \times .

retinal transcripts of dog or mouse by RT-PCR or 5' RACE. The correct sequence was confirmed, and the final ORF was chosen based on conservation between species. *PRCD* encodes a 54-aa protein in dog and human; in mouse it is 53 aa as 3 in-frame coding nucleotides (Nos. 28–30) are missing from exon 1. The conservation of potential regulatory elements and binding sites further suggests the correct positioning of the proposed ORF (Supplementary Fig. 1). Three CRX binding sites are located just upstream of the ATG site in dog, human, and mouse; this factor is thought to drive photoreceptor expression [23,24]. A large number of potential transcription factor binding sites could be identified within the 5'UTR and intron 1, and 13 of these were common to the three species investigated. However, only the Sox17 site showed high positional conservation, suggesting that this might be one of the *PRCD*-interacting genes even though this transcription factor predominantly regulates the transcription of endodermal genes [25,26]. Additional studies currently are under way to assess further the regulation and function of *PRCD*.

Comparison of the cDNA between *prcd*-affected and normal dogs identified a G to A substitution at position 5 of the canine *PRCD* ORF that causes a C2Y change in the translated protein. No other change within the available RPCI81 BAC 10-M13 sequence was found that segregated with the disease phenotype. Subsequent screening of the canine *PRCD* mutation indicated the absence of the mutated allele (A) in 1120 dogs from 91 different breeds not diagnosed with the *prcd* form of PRA as well as 24 red and 6 gray wolves (Supplementary Tables 2 and 3). As well, breeds identified to segregate the *prcd* haplotype [16] showed complete concordance between the mutated allele in the homozygous state and the clinical diagnosis of *prcd*. The high number of dogs investigated supports the association between mutation and disease phenotype, and this is particu-

larly important when identifying a mutation in a novel gene of unknown function.

Some exceptions in the investigated breeds, however, were noted and bear mentioning. We have identified dogs with retinal degeneration that were homozygous for the normal (G) allele. Labrador retrievers with advanced retinal degeneration at <1 year rather than 6–8 years represented age outliers for the breed-specific disease [2]. Others, e.g., CC, ST, and ACD, had retinal degeneration that was clinically similar to *prcd*, either in age of onset or in clinical appearance, but were homozygous normal for canine *PRCD*. It is likely that these retinal degenerations, if not acquired phenocopies, are caused by mutations in different retinal disease genes, an issue that we have previously reported for the Norwegian elkhound and miniature schnauzer breeds (see [2] for review). For all of the discordant individuals for which haplotype analysis was carried out, we were able to exclude the segregating chromosomes from association with *prcd* [16].

An AE dog was of particular interest because she typed homozygous affected for *prcd* (A/A), but did not show clinical signs at 13 years. Both chromosomes were confirmed to produce the disease phenotype in affected relatives through extended haplotype analysis. Such a delay in onset of disease is not surprising given the variation in disease onset and progression that occurs in some breeds with *prcd* [9]. For example, in the ECS we have found the age of early diagnosis to vary from 3.1 to 13.5 years in dogs having the same mutation in canine *PRCD* (OptiGen LLC, unpublished information). Comparative analysis of the genetic background between different phenotypic variants of *prcd* has been proposed to identify the genetic modifiers. Such studies will contribute to our understanding of *PRCD* function and the molecular mechanisms of disease.

The cloning and characterization of the human *PRCD* ortholog, and the subsequent screening of patients clinically diagnosed with various forms of retinal disease, led to the identification of a patient that is homozygous for the same sequence variation, at both the nucleotide and the amino acid level, as found in *prcd* dogs. This coincidence may suggest that function-altering sequence variations that occur elsewhere in this gene are likely to have a phenotype different from that of the patients that were screened in this study. The one RP patient identified with a *PRCD* mutation had advanced disease in the fourth decade of life with marked constriction of the visual fields, a nonrecordable electroretinogram, and extensive bone-spicule-like pigmentation. Such advanced disease is similar to what is observed in most canines affected with *prcd* at a comparable age [2]. Although the patient has a family history of vision problems, the remaining family members still live in Bangladesh, and the acquisition of additional samples and phenotype characterization of family members has been difficult. We are hopeful these data will become available soon. Although this study shows that human *PRCD* mutations are an extremely rare cause of retinitis pigmentosa in patients ascertained in North America, it will require further study to determine whether mutations in this gene are responsible for a larger fraction of this disease in Bangladesh or in other Asian populations.

Expression of *PRCD* indicates two main retinal transcripts of 2 and 6.5 kb using northern analysis, both of which contain exons 1–4, as well as the originally identified 3'UTR region. The expression of canine *PRCD* was determined by qRT-PCR and showed that the highest level is in retina followed by RPE/choroid and that other tissues have very low or negligible levels. However, the use of laser microscopy to isolate specifically different retinal cell layers allowed us to quantify more precisely the contribution of each to the total canine *PRCD* mRNA pool. This showed that there was approximately equal expression in the RPE, photoreceptor, and ganglion cell layers, thus explaining why there is no decrease in canine *PRCD* expression as the photoreceptor cells are lost with disease progression. Data from immunocytochemical subcellular localization experiments suggest that the mutant form of the protein exhibits a distinct pattern of localization into spherical profiles within the cytoplasm, and we posit that this altered localization may impair function and/or its interactions with other proteins in the affected human and canine retina.

As can be appreciated, it is extremely difficult to assess the function of a protein that has not been known thus far and does not belong to any known gene family. To date, we have shown that *PRCD* is causally associated with retinal degeneration in dog and human. As well, *in silico* analysis strongly suggests the functional importance of this gene based on the high conservation in vertebrate species. Our working hypothesis includes the formation of an α -helix involving the first 24 aa of *PRCD* that might contain a transmembrane segment; partial homology of this region to other confirmed or predicted bacterial proteins (e.g., AAS62120, ABD02230) highly supports this hypothesis. The first 4 aa, totally conserved in 14 vertebrate species analyzed, are not predicted to be part of the transmembrane domain. They are likely to play an important

role as the recognition site for interacting proteins. Therefore, the described mutation in the highly conserved second amino acid would significantly influence the function of *PRCD*. The nature of the mutation and the observed change in the localization are concordant with changes in the long-term viability of the retina rather than a fast-acting developmental disorder. However, the exact disease mechanism will require a series of further studies.

The importance of this novel gene, *PRCD*, for maintenance of rod photoreceptor structure, function, and/or survival is attested to by its high level of conservation in vertebrate species. The significance of this conservation, absolute for the first four codons, is underscored by the causal association of the C2T mutation reported here with hereditary retinal degeneration in both dogs and humans. The apparent absence of a homologous gene in invertebrates suggests that the function of the *PRCD* protein is critically important to those aspects of photoreceptor structure or metabolism unique to vertebrate visual cells. As these issues yield to further studies, they should provide new insights into vertebrate photoreceptor evolution, as well as into the mechanisms of photoreceptor health and disease.

Materials and methods

FISH mapping

Four milliliters of peripheral blood containing 0.2 ml preservative-free heparin was incubated with 31 ml of complete medium for 96 h at 37°C [27]. After incubation, 5 ml of medium was aspirated, and cells were arrested with 0.2 ml colcemid (10 μ /ml) during a 1.5-h incubation. Chromosomal spreads were prepared on slides using standard hypotonic solution and methanol/acetic acid fixation, and chromosome painting was performed [28] using RPC181 BAC clones specific to CFA9 (No. 189-B1) or included in the physical map (No. 429-O8) of the *prcd* interval [15]. FISH images of probes hybridized onto G-banded metaphases were captured using the ISIS software (In Situ Imaging System; Metasystems) with a Paco CCD camera mounted on an Aristoplan (Leitz) microscope [28].

Protein structural analysis

Three-dimensional (3D) models for *PRCD* sequences were built using the MODELLER program [29–31] with (a) experimentally determined structures for the templates extracted from the Protein Data Bank and (b) sequence alignments obtained from the BIOINFO metaserver [32]. The pair-wise alignments produced by the metaserver were assigned relatively low scores (low level of certainty) by the 3-D-Jury tool. However, the set of highest score alignments points to structures having a short α -helical fragment at the N-terminal end of the protein. 3D models for all the sequences were generated by comparative protein modeling through satisfaction of spatial constraints with the MODELLER program, minimizing infringements of distance and dihedral-angle restraints from the template structures.

Study samples

Research dogs were maintained at the Retinal Disease Studies Facility in Kennett Square, Pennsylvania, under cyclic light conditions, and *prcd* status was established using morphological, ERG, or genetic criteria [14]. The dogs represented several breeds in which *prcd* allelism had been established through interbreed crosses [2,9]: ECS, LR, MP, and TP. Allelism with *prcd* has also been established for ACD, NSDTR, and PWD [16]. In addition to the colony-derived research dogs, we obtained blood samples and clinical records of dogs submitted by their owners or other scientists directly to our lab or through OptiGen, LLC.

In all animals maintained in the research colony, and many outside cases, two of the investigators (G.M.A., G.D.A.) carried out the clinical evaluations.

Euthanasia with a barbiturate overdose was performed at a single time of the day (10:00 AM) to prevent fluctuation in the abundance of retinally expressed genes, and retinas were isolated from the eyecup within 1–2 min of death. Other tissues were obtained within 5–10 min, and all tissues were flash frozen in liquid nitrogen and stored at -80°C until used. The human retina was isolated from a 76-year-old female donor 8 h postmortem and obtained through the National Disease Research Interchange. After isolation, it was snap frozen in liquid nitrogen and processed as described. Neural retina from C57BL/6 mice were pooled to obtain sufficient tissue for cDNA amplification of the murine *Prcd*.

A total of 1863 unrelated human patients with a clinical diagnosis of heritable photoreceptor disease were studied. All individuals provided written informed consent. Approximately a third of these patients were ascertained in the Retina Clinic of the University of Iowa, while the remainder was ascertained by inherited retinal disease specialists elsewhere in the United States. One thousand two hundred forty-one patients were diagnosed with RP, 283 with Leber congenital amaurosis, 224 with cone-rod dystrophy, and 115 with Usher syndrome. All of these patients had been screened previously for mutations in one or more genes known to cause their diseases (RetNet—Retinal Information System; <http://www.sph.uth.tmc.edu/RetNet/>), and none of them had been found to harbor a plausible disease-causing variation in a known retinal disease gene. Normal controls from the University of Iowa consisted of 166 subjects (general population controls) of mixed ages who had no personal or family history of heritable retinal disease. Over 80% of this group described themselves as “Caucasian.” The second group consisted of 49 unrelated control subjects from India and Pakistan.

Sample preparation

Total RNA was isolated using Trizol extraction followed by treatment with DNase (1 unit RQ1 DNase; Promega), if used for qRT-PCR, and protection with RNase inhibitor (Roche Applied Science). Double-stranded cDNA was synthesized using the Thermoscript RT-PCR kit (Invitrogen) with random hexamer primers. Amplification was obtained under standard conditions (Supplementary Table 1, primers 1, 2, and 11; Accession Nos. DQ390332, DQ390333), and extension of cDNA sequence by RACE was obtained from a 10-week normal retina through the SMARTTRACE cDNA amplification kit (Clontech) following the manufacturer’s protocol (Supplementary Table 1, primers 5–8). RT-PCR using primers from exons 1 and 4 (Supplementary Table 1, primers 9 and 10) confirmed the main transcript (695 bp, four exons; Accession No. DQ390330), but suggested a second transcript, which does not splice out intron 1 (Accession No. DQ390331). Additionally, primers located specifically in exon 2 and the predicted exon 5 of canine *PRCD* (Supplementary Table 1, primers 12 and 13) yielded three splice variants (796, 763, and 509 bp), utilizing different potential splicing sites between exons 4 and 5 (Accession Nos. DQ390334, DQ390335, DQ390336). All amplicons were cloned into the pCR 2.1-TOPO vector (Invitrogen) following the manufacturer’s protocol and stored in glycerol. Selected clones were sequenced for insert verification. DNA was extracted from spleen or blood using the QIAmp DNA Mini Kit (Qiagen). All RNA and DNA samples were evaluated on a 1% denaturing agarose–formaldehyde or agarose gel prior to use, and level of purity and concentration was measured with a spectrophotometer.

Restriction enzyme analysis

Amplification from genomic DNA with primers 3 and 17 (Supplementary Table 1) at an annealing temperature of 58°C under conditions described above resulted in a 512-bp PCR product. Restriction enzyme digestion performed in a 25- μl reaction on 500 ng PCR product with *RsaI* resulted in 396- and 116-bp bands in the presence of the mutated (A) allele; digestion with *ApaI* resulted in 397- and 115-bp bands in the presence of the normal (G) allele. The genotypes resulting from the digestion were visualized on 6% polyacrylamide gels.

Pyrosequencing

High-throughput screening for the mutant (A) and normal (G) *PRCD* alleles was done by Pyrosequencing (Biotage). PCR was performed on isolated DNA

with primers (Supplementary Table 1; primer 18 and biotin-labeled primer 19) under standard conditions with 1.5 mM MgCl_2 at 55°C annealing temperature in 35 cycles, checked on a 2% agarose gel, and purified with Streptavidin Sepharose High Performance Beads (Amersham). Primer 20 was added to initiate the reaction, and primer elongation was achieved and recorded on a Pyrosequencer PSQ 96MA using the PSQ 96 SNP reagent kit (Biotage).

Human *PRCD* screen

DNA was extracted from peripheral blood according to a previously described protocol [33]. Samples from 1863 patients and 215 controls were screened for variations in the whole coding sequence of *PRCD* using single-strand conformational polymorphism (SSCP) analysis as previously described [34]. To conserve DNA, the 49 Indian/Pakistani control samples were screened for variations in exon 1 only. All samples exhibiting an abnormal electrophoretic pattern by SSCP were further characterized using automated bidirectional sequencing on an ABI Model 3730 using the BigDye version 3 chemistry (Applied Biosystems).

Northern analysis

Samples were loaded on a 1% agarose–formaldehyde denaturing gel, and 3 μg of 0.24–9.5 kb RNA ladder was used as a size marker (Invitrogen). The gel ran with continuously circulating $1\times$ Mops running buffer (Ambion) for 16 h at 21 V. After three 5-min washes in DEPC-treated water, 20 min in 0.05 N NaOH, and a 15-min soak in $10\times$ SSC, transfer to a nylon-based membrane (GeneScreen Plus, NEN Life Science) was done with $10\times$ SSC buffer using a standard protocol. Full transfer was confirmed by exposing the gel to UV light. The membrane was washed in $2\times$ SSC for 2 min, and RNA was cross-linked to the membrane (exposure was 0.12 J/cm²; Stratilinker UV Crosslinker).

Northern probes were amplified from cDNA clones and labeled with [α -³²P]dCTP using the RadPrime DNA labeling system (Invitrogen). Prehybridization and hybridization were carried out with ExpressHyb solution (Clontech) at 68°C for 30 min. Hybridization was carried out at 68°C for 16–18 h, and the blots were rinsed several times with $2\times$ SSC, 0.05% SDS; the washes with the same solution were done twice with continuous agitation for 40 min. Then the blot was washed with $0.1\times$ SSC and 0.1% SDS with continuous shaking at 50°C for 40 min with one change of fresh solution. Blots were exposed to X-ray film at and 70°C for 24–96 h with two intensifying screens. Loading control was achieved by hybridizing a canine-specific β -actin (Z70044) probe to the membranes under the same conditions and exposure to X-ray film for 4 h.

Quantitative real time PCR

Expression profiling of *PRCD* was done on a 7500 Real Time PCR System (Applied Biosystems) with two TaqMan assays. Probes and primers were designed using the Primer Express Software (Applied Biosystems) resulting in two primer sets. One produced a 55-bp product specific for exon 1 (Supplementary Table 1, primers 24 and 25, probe 26) and the second a 63-bp product spanning the exon 2 and 3 junction (Supplementary Table 1, primers 21 and 22, probe 23). TaqMan PCR amplification was performed in a two-step RT-PCR, reverse transcribing the RNA first with the TaqMan reverse transcription reagents (Applied Biosystems) followed by amplification of the target sequence using the TaqMan Universal Master Mix with the specific probe sets. Gene expression was evaluated through relative quantification against an 18S endogenous control (Hs99999901_s1; Applied Biosystems).

Laser capture microscopy

Glass slides with 10- μm frozen sections of normal canine retina (16 weeks) fixed in 4% paraformaldehyde and embedded in OCT [35] were thawed for 20 s and subsequently dehydrated through serial immersion in 70, 95, and 100% EtOH for 1 min each, followed by 2×5 min in xylene. Laser capture was performed with a PixCell II laser capture microdissection apparatus (Arcturus) on CapSure HS LCM Caps (Arcturus) at 50 mW, capturing a total of 3024 ganglion, 2525 RPE, and 1850 photoreceptor shots. RNA was isolated with the

PicoPure RNA isolation kit (Arcturus) and cDNA amplified with the RiboAmp OA RNA amplification kit (Arcturus), both following manufacturer's protocols. A final cDNA concentration of ~15 ng/μl was obtained based on spectrophotometer measurement for each cell class isolated, and 1 μl was used for qRT-PCR as described.

Protein expression and localization

COS-7 cells (American Type Culture Collection, Manassas, VA, USA) were maintained in Dulbecco's modified Eagle's medium [with 10% (v/v) fetal calf serum, 1% penicillin/streptomycin, L-glutamine, sodium pyruvate, nonessential amino acids, Hepes, and gentamycin]. Prior to transfection, cells were split and grown on coverslips in six-well plates. Transfection of expression vectors (pCDNA3.1v5His; see above) containing either the wild-type human *PRCD* gene or the C2Y mutant were mediated using Lipofectamine 2000 (Invitrogen) in Opti-MEM 1 (Gibco) for 48 h according to the manufacturer's instructions.

Cells were fixed for 5 min in 4% formaldehyde (prepared from paraformaldehyde) in 10 mM phosphate-buffered saline (PBS), pH 7.4. Coverslips were then washed in PBS, and their top surfaces were covered with blocking solution (1 mg/ml bovine serum albumin in PBS) for 15 min. Following a brief rinsing step in PBS, coverslips were then incubated with anti-His tag primary antibody (Invitrogen) at a concentration of 6 μg/ml in PBS with 0.1% Triton X-100 for 1 h. After being rinsed with PBS, coverslips were incubated with an anti-mouse secondary antibody conjugated to Alexa-546 (Invitrogen) at a 1 to 200 dilution for 30 min. Coverslips were washed and then counterstained with 4',6-diamidino-2-phenylindole (DAPI) and with Alexa 488-conjugated phalloidin, an actin-binding dye, diluted 1 to 200 (Invitrogen). Coverslips were then placed on glass slides, mounted with Aquamount (Ted Pella), and photographed with a BX41 fluorescence microscope (Olympus).

Acknowledgments

We are grateful to Dr. Charles Aquadro, Department of Molecular Biology and Genetics, Cornell University, for help with pyrosequencing; Dr. Peter Felsburg in the Section of Medical Genetics, University of Pennsylvania, for access to the Applied Biosystems 7500 Real-Time PCR System; and Dr. Neal Rubinstein, Department of Cell and Developmental Biology and Pennsylvania Muscle Institute, University of Pennsylvania, for access to and help with laser capture microscopy. Some blood samples with pedigrees and clinical reports were provided by Dr. David Sargan (FL), Dr. Keith Murphy (AE subset), Dr. Simon Petersen-Jones (EMD), The Seeing Eye, Inc. (LR subset), Dr. Ron Riis, Dr. Elaine Ostrander (many nonaffected breeds), and many interested dog breeders and breed clubs. Dr. Jean Bennett provided normal mouse retinas for RNA extraction, and Ulana Prociuk assisted with chromosomal spreads. The authors especially acknowledge Jennifer Johnson, Julie Jordan, and Amy Antosh for technical assistance; Jean Andorf and Louisa Affatigato for sequencing of human samples; Drs. Jaroslaw Pillardy and Qi Sun for bioinformatics support; and Amanda Nickle, Gerri Antonini, and staff of the RDS facility for care and supervision of the research dog colony. This article is dedicated to Sue Van Sloun for steadfast support, encouragement, and help. This work was supported by the Foundation Fighting Blindness, the Morris Animal Foundation/The Seeing Eye, Inc., NIH Grants EY06855 and EY13132, the Van Sloun Fund for Canine Genetic Research, the L.J. Niles Foundation, the ONCE International Prize for Biomedicine and R&D for New Technologies for the Blind, targeted donations from many

individuals and breed clubs, and the HHMI. The discovery of the *PRCD* gene and potential tests identifying the mutation in the dog have been patented. Authors B. Zangerl, O. Goldstein, S.E. Pearce-Kelling, J.S. Felix, G.M. Acland, and G.D. Aguirre hold the rights to this patent through Cornell University, and authors J.S. Felix, G.M. Acland, and G.D. Aguirre are co-owners of the company (OptiGen, LLC, Ithaca, NY, USA) that has licensed the technology from Cornell University.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.ygeno.2006.07.007](https://doi.org/10.1016/j.ygeno.2006.07.007).

References

- [1] G. Aguirre, J. Alligood, P. O'Brien, N. Buyukmihci, Pathogenesis of progressive rod–cone degeneration in miniature poodles, *Invest. Ophthalmol. Visual Sci.* 23 (1982) 610–630.
- [2] G.D. Aguirre, G.M. Acland, Models, mutants and man: searching for unique phenotypes and genes in the dog model of inherited retinal degeneration, in: E.A. Ostrander, U. Giger, K. Lindblad-Toh (Eds.), *The Dog and Its Genome*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 2006, pp. 291–325.
- [3] B. Wiggert, G. Kutty, K.O. Long, L. Inouye, I. Gery, G.J. Chader, G.D. Aguirre, Interphotoreceptor retinoid-binding protein (IRBP) in progressive rod–cone degeneration (*prcd*)—Biochemical, immunocytochemical and immunologic studies, *Exp. Eye Res.* 53 (1991) 389–398.
- [4] R.E. Anderson, M.B. Maude, G.M. Acland, G.D. Aguirre, Plasma lipid changes in *PRCD*-affected and normal miniature poodles given oral supplements of linseed oil: indications for the involvement of n-3 fatty acids in inherited retinal degenerations, *Exp. Eye Res.* 58 (1994) 129–137.
- [5] K. Miezewska, T. van Veen, G.D. Aguirre, Structural changes of the interphotoreceptor matrix in an inherited retinal degeneration: a lectin cytochemical study of progressive rod–cone degeneration, *Invest. Ophthalmol. Visual Sci.* 34 (1993) 3056–3067.
- [6] G.D. Aguirre, L. Andrews, Nomarski evaluation of rod outer segment renewal in a hereditary retinal degeneration: comparison with autoradiographic evaluation, *Invest. Ophthalmol. Visual Sci.* 28 (1987) 1049–1058.
- [7] G. Aguirre, P. O'Brien, Morphological and biochemical studies of canine progressive rod–cone degeneration: ³H-fucose autoradiography, *Invest. Ophthalmol. Visual Sci.* 27 (1986) 635–655.
- [8] J.H. Parkes, G.D. Aguirre, J.H. Rockey, P.A. Liebman, Progressive rod–cone degeneration in the dog: characterization of the visual pigment, *Invest. Ophthalmol. Visual Sci.* 23 (1982) 674–678.
- [9] G.D. Aguirre, G.M. Acland, Variation in retinal degeneration phenotype inherited at the *prcd* locus, *Exp. Eye Res.* 46 (1988) 663–687.
- [10] J.C. Huang, M.F. Chesselet, G.D. Aguirre, Decreased opsin mRNA and immunoreactivity in progressive rod–cone degeneration (*prcd*): cytochemical studies of early disease and degeneration, *Exp. Eye Res.* 58 (1994) 17–30.
- [11] G.D. Aguirre, G.M. Acland, M.B. Maude, R.E. Anderson, Diets enriched in docosahexaenoic acid fail to correct progressive rod–cone degeneration (*prcd*) phenotype, *Invest. Ophthalmol. Visual Sci.* 38 (1997) 2387–2407.
- [12] K.E. Gropp, J.C. Huang, G.D. Aguirre, Differential expression of photoreceptor-specific proteins during disease and degeneration in the progressive rod–cone degeneration (*prcd*) retina, *Exp. Eye Res.* 64 (1997) 875–886.
- [13] S. Bardiën, R. Ramesar, S. Bhattacharya, J. Greenberg, Retinitis pigmentosa locus on 17q (RP17): fine localization to 17q22 and exclusion of the *PDEG* and *TIMP2* genes, *Hum. Genet.* 101 (1997) 13–17.
- [14] G.M. Acland, K. Ray, C.S. Mellersh, W. Gu, A.A. Langston, J. Rine, E.A. Ostrander, G.D. Aguirre, Linkage analysis and comparative mapping of

- canine progressive rod–cone degeneration (prcd) establishes potential locus homology with retinitis pigmentosa (RP17) in humans, *Proc. Natl. Acad. Sci. USA* 95 (1998) 3048–3053.
- [15] D.J. Sidjanin, B. Miller, J. Kijas, J. McElwee, J. Pillardy, J. Malek, G. Pai, T. Feldblyum, C. Fraser, G.M. Acland, G.D. Aguirre, Radiation hybrid map, physical map, and low-pass genomic sequence of the canine prcd region on CFA9 and comparative mapping with the syntenic region on human chromosome 17, *Genomics* 81 (2003) 138–148.
- [16] O. Goldstein, B. Zangerl, S.E. Pearce-Kelling, D.J. Sidjanin, J.W. Kijas, J.S. Felix, G.M. Acland, G.D. Aguirre, Linkage disequilibrium mapping in domestic dog breeds narrows the progressive rod–cone degeneration interval and identifies ancestral disease-transmitting chromosome, *Genomics* 88 (2006) 541–550.
- [17] R. Li, E. Mignot, J. Faraco, H. Kadotani, J. Cantanese, B. Zhao, X. Lin, L. Hinton, E.A. Ostrander, D.F. Patterson, J.P. de Jong, Construction and characterization of an eightfold redundant dog genomic bacterial artificial chromosome library, *Genomics* 58 (1999) 9–17.
- [18] B. Zangerl, Q. Sun, J. Pillardy, J.L. Johnson, P.A. Schweitzer, A.G. Hernandez, L. Liu, G.M. Acland, G.D. Aguirre, Development and characterization of a normalized canine retinal cDNA library for genomic and expression studies, *Invest. Ophthalmol. Visual Sci.* 47 (2006) 2632–2638.
- [19] K. Lindblad-Toh, C.M. Wade, T.S. Mikkelsen, E.K. Karlsson, D.B. Jaffe, M. Kamal, M. Clamp, J.L. Chang, E.J. Kulbokas, M.C. Zody, E. Mauceli, X. Xie, M. Breen, R.K. Wayne, E.A. Ostrander, C.P. Ponting, F. Galibert, D.R. Smith, P.J. Dejong, E. Kirkness, P. Alvarez, T. Biagi, W. Brockman, J. Butler, C.W. Chin, A. Cook, J. Cuff, M.J. Daly, D. Decaprio, S. Gnerre, M. Grabherr, M. Kellis, M. Kleber, C. Bardeleben, L. Goodstadt, A. Heger, C. Hitte, L. Kim, K.P. Koepfli, H.G. Parker, J.P. Pollinger, S.M. Searle, N.B. Sutter, R. Thomas, C. Webber, J. Baldwin, A. Abebe, A. Abouelleil, L. Aftuck, M. Ait-Zahra, T. Aldredge, N. Allen, P. An, S. Anderson, C. Antoine, H. Arachchi, A. Aslam, L. Ayotte, P. Bachantsang, A. Barry, T. Bayul, M. Benamara, A. Berlin, D. Bessette, B. Blitshteyn, T. Bloom, J. Blye, L. Boguslavskiy, C. Bonnet, B. Boukhgalter, A. Brown, P. Cahill, N. Calixte, J. Camarata, Y. Cheshatsang, J. Chu, M. Citroen, A. Collymore, P. Cooke, T. Dawoe, R. Daza, K. Decktor, S. Degray, N. Dhargay, K. Dooley, K. Dooley, P. Dorje, K. Dorjee, L. Dorris, N. Duffey, A. Dupes, O. Egbiremolan, R. Elong, J. Falk, A. Farina, S. Faro, D. Ferguson, P. Ferreira, S. Fisher, M. Fitzgerald, et al., Genome sequence, comparative analysis and haplotype structure of the domestic dog, *Nature* 438 (2005) 803–819.
- [20] W. Makalowski, M.S. Boguski, Evolutionary parameters of the transcribed mammalian genome: an analysis of 2,820 orthologous rodent and human sequences, *Proc. Natl. Acad. Sci. USA* 95 (1998) 9407–9412.
- [21] J.A. Moody, T.R. Famula, R.C. Sampson, K.E. Murphy, Identification of microsatellite markers linked to progressive retinal atrophy in American Eskimo dogs, *Am. J. Vet. Res.* 66 (2005) 1900–1902.
- [22] M. Heitmann, H. Hamann, R. Brahm, H. Grussendorf, C.U. Rosenhagen, O. Distl, Analysis of prevalence of presumed inherited eye diseases in Entlebucher mountain dogs, *Vet. Ophthalmol.* 8 (2005) 145–151.
- [23] T. Furukawa, E.M. Morrow, C.L. Cepko, Crx, a novel otx-like homeobox gene, shows photoreceptor-specific expression and regulates photoreceptor differentiation, *Cell* 91 (1997) 531–541.
- [24] T. Furukawa, E.M. Morrow, T. Li, F.C. Davis, C.L. Cepko, Retinopathy and attenuated circadian entrainment in Crx-deficient mice, *Nat. Genet.* 23 (1999) 466–470.
- [25] D. Sinner, S. Rankin, M. Lee, A.M. Zorn, Sox17 and beta-catenin cooperate to regulate the transcription of endodermal genes, *Development* 131 (2004) 3069–3080.
- [26] T. Niimi, Y. Hayashi, S. Futaki, K. Sekiguchi, SOX7 and SOX17 regulate the parietal endoderm-specific enhancer activity of mouse laminin alpha1 gene, *J. Biol. Chem.* 279 (2004) 38055–38061.
- [27] P. Werner, M.G. Raducha, U. Prociuk, P.S. Henthorn, D.F. Patterson, Physical and linkage mapping of human chromosome 17 loci to dog chromosomes 9 and 5, *Genomics* 42 (1997) 74–82.
- [28] A.S. Graphodatsky, F. Yang, P.C. O'Brien, N. Serdukova, B.S. Milne, V. Trifonov, M.A. Ferguson-Smith, A comparative chromosome map of the Arctic fox, red fox and dog defined by chromosome painting and high resolution G-banding, *Chromosome Res.* 8 (2000) 253–263.
- [29] R. Sanchez, A. Sali, Comparative protein structure modeling: introduction and practical examples with modeller, *Methods Mol. Biol.* 143 (2000) 97–129.
- [30] A. Sali, L. Potterton, F. Yuan, H. van Vlijmen, M. Karplus, Evaluation of comparative protein modeling by MODELLER, *Proteins* 23 (1995) 318–326.
- [31] A. Sali, T.L. Blundell, Comparative protein modelling by satisfaction of spatial restraints, *J. Mol. Biol.* 234 (1993) 779–815.
- [32] K. Ginalski, A. Elofsson, D. Fischer, L. Rychlewski, 3D-Jury: a simple approach to improve protein structure predictions, *Bioinformatics* 19 (2003) 1015–1018.
- [33] G.J. Buffone, G.J. Darlington, Isolation of DNA from biological specimens without extraction with phenol, *Clin. Chem.* 31 (1985) 164–165.
- [34] A.R. Webster, E. Heon, A.J. Lotery, K. Vandenberg, T.L. Casavant, K.T. Oh, G. Beck, G.A. Fishman, B.L. Lam, A. Levin, J.R. Heckenlively, S.G. Jacobson, R.G. Weleber, V.C. Sheffield, E.M. Stone, An analysis of allelic variation in the ABCA4 gene, *Invest. Ophthalmol. Visual Sci.* 42 (2001) 1179–1189.
- [35] W.A. Beltran, H. Rohrer, G.D. Aguirre, Immunolocalization of ciliary neurotrophic factor receptor alpha (cntfralpha) in mammalian photoreceptor cells, *Mol. Vision* 11 (2005) 232–244.